

Raised cholesterol and breastfeeding

The information provided is taken from various reference sources. It is provided as a guideline. No responsibility can be taken by the author or the Breastfeeding Network for the way in which the information is used. Clinical decisions remain the responsibility of medical and breastfeeding practitioners. The data presented here is intended to provide some immediate information but cannot replace input from professionals including <https://www.sps.nhs.uk/>.

If a breastfeeding mother has raised cholesterol (not familial) encourage diet and lifestyle changes and monitor at intervals using the Qrisk calculator <https://qrisk.org/three/>

The treatment of Familial Hypercholesterolemia is recommended to reduce the risk of future cardiovascular events. The balance is between the treatment of the breastfeeding mother with medication (on which there are few studies) and the ongoing value of breastfeeding which has cardio-protective advantages for mother and baby.

1. If the breastfeeding mother has Familial Hypercholesterolemia, it is assumed that even reducing the level in maternal milk will still exceed that in standard infant formula together with added protective cardio-vascular properties of breastmilk (Holmsen)
2. Standard Infant formula contains no cholesterol. Cholesterol is essential for healthy development of the baby. (Lawrence) and has none of the other positive benefits of breastfeeding which is the biological norm.
3. Breastfeeding has many factors to protect the mother and baby from future cardiovascular disease.

NB This data has been taken from expert sources but there is currently an absence of data on the effect of statins on breastfed babies.

This document provides an overview of research and is not intended as a recommendation. Treatment of raised cholesterol in breastfeeding should be made after informed discussion between parents and their health professionals whilst protecting breastfeeding.

The first-line choice of cholesterol-lowering treatment for breastfeeding parents who do not have Familial Hypercholesterolemia is Cholestyramine (<https://www.ncbi.nlm.nih.gov/books/NBK501394/>). It is compatible with breastfeeding. This factsheet has been developed to provide information on "statins" as well as some of the newer therapies.

Cholesterol

Cholesterol is necessary for the development of brain tissue, myelination of nerves, and is the basis for many enzymes. Breastfed infants have higher plasma cholesterol levels than those fed standard artificial formulas as these products contain no cholesterol at all. The higher cholesterol levels in breastmilk protects babies against the consequences of hypercholesterolemia in adult life (Lawrence 2016).

To talk to a mum who knows about breastfeeding call the National Breastfeeding Helpline 0300 100 0212

Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package.

Cholesterol tests measure (NHS):

- total cholesterol – the overall amount of cholesterol in your blood, including both "good" and "bad" cholesterol
- good cholesterol (called HDL) – this makes you less likely to have heart problems or a stroke
- bad cholesterol (called LDL and non-HDL) – this makes you more likely to have heart problems or a stroke
- triglycerides – a fatty substance similar to bad cholesterol

Breastfeeding and Cardio-vascular disease (CVD)

Research shows that being breastfed leads to better outcomes with respect to cardiovascular disease in later life (UNICEF). Cessation of breastfeeding may have consequences for the mother and baby and a recommendation to stop should not be undertaken without examining the literature for benefit and risk to the baby of the medication.

Nguyen (2019) reported that ever breastfeeding was associated with lower risk of CVD hospitalisation and mortality compared with never breastfeeding, and breastfeeding ≤ 12 months/child was significantly associated with lower risk of CVD hospitalisation. WHO (2021) and UNICEF recommend: early initiation of breastfeeding within 1 hour of birth; exclusive breastfeeding for the first 6 months of life; and introduction of nutritionally adequate and safe complementary (solid) foods at 6 months together with continued breastfeeding up to 2 years of age or beyond.

Schwartz (2009) studied 139,681 women who had breastfed for more than 12 months across their lactations and showed a 10-15% reduction in hypertension, diabetes, hyperlipidaemia, and cardiovascular disease than those who had never breastfed.

Hui (2019) looked at an association between breastfeeding in the first three months of life with lipid profile and adiposity at around 17.5 years in a population in Hong Kong. The team found that exclusive breastfeeding, (but not mixed feeding) at 0 to 3 months, compared with artificial formula feeding was associated with lower total cholesterol and low-density lipoprotein (LDL) cholesterol but not with high-density lipoprotein cholesterol (HDL). LDL is sometimes referred to as "bad" cholesterol because it collects in the walls of blood vessels. HDL is seen as "good" cholesterol, because it absorbs cholesterol and carries it back to the liver. The liver then flushes it from the body. High levels of HDL cholesterol can lower the risk for heart disease and stroke.

Singhal (2004) studied 926 preterm babies and determined evidence for the long-term benefits of breastmilk feeding on the risk of atherosclerosis.

Owen (2002) conducted a cross-sectional study of 1,532 adolescents in 10 British towns and determined that breastfeeding is associated with increased mean serum total cholesterol and low-density lipoprotein cholesterol in infancy but with lower levels in adult life providing long-term benefits for cardiovascular health.

These studies should not be taken as the advantages of breastfeeding but rather the risk of deleterious health outcomes for babies exclusively fed with infant formula (Renfrew 2012).

Raised cholesterol

Cholesterol is a fatty substance which is made in the liver. It is also found in foods. The Mediterranean diet is generally recommended to decrease cholesterol obtained through consumption of foods.

Cholesterol levels may be found to be raised at routine monitoring of a woman. Diet and lifestyle issues should be addressed before considering the initiation of cholesterol lowering medication to avoid unnecessary medication. Raised cholesterol is mainly caused by:

- eating fatty food,
- not exercising enough,
- being overweight,
- smoking and drinking alcohol

It is recommended that those with high cholesterol eat more:

- oily fish, like mackerel and salmon
- brown rice, bread, and pasta
- nuts and seeds
- fruits and vegetables

and eat less:

- meat pies, sausages, and fatty meat
- butter, lard, and ghee
- cream and hard cheese, like cheddar
- cakes and biscuits
- food that contains coconut oil or palm oil (NHS)

They should aim to exercise more (150 minutes (2.5 hours)) a week. This might include walking until the heart starts beating faster (pushing a pram round a local park or carrying baby in a sling whilst you walk), swimming or cycling. However, as a new mother it is acknowledged this can be difficult. Smoking cessation can improve the levels too (for more information:

<https://www.breastfeedingnetwork.org.uk/smoking/>)

When undertaking a review of cardiovascular risk, a calculator is used to determine the likelihood of cardiovascular disease occurring in the next 10 years. Risk calculators should not be used for people already identified as being at high risk, such as those with diabetes or Familial Hypercholesterolaemia (see below). The risk is calculated considering many other factors including height, weight, age, smoking status, alcohol consumption, blood pressure, family history, chronic medical conditions, cholesterol level etc (<https://qrisk.org/three/>). This permits discussion with a health professional on possible lifestyle changes or initiation of medication. (See [Your NHS Health Check results and action plan - NHS \(www.nhs.uk\)](#) for further information.)

Cholestyramine may be considered as first line if medication continues to be necessary due to its low oral bioavailability (0%) so that so it cannot transfer to breastmilk, nor to the infant's plasma via breastmilk (<https://www.ncbi.nlm.nih.gov/books/NBK501394/>).

Familial Hypercholesterolaemia

Women with Familial Hypercholesterolaemia are currently advised to discontinue breastfeeding prior to beginning statin therapy after pregnancy. This is because we have little data from studies on the compatibility of statins with breastfeeding. There has always been a concern that we do not know what effect potentially lowering cholesterol in the breastfed baby has.

Familial Hypercholesterolaemia (FH) is an inherited condition which can lead to extremely high cholesterol levels. Without treatment, FH can lead to heart disease at a very young age. But once it's been diagnosed, it can be treated with medicines and a healthy lifestyle (HEARTUK). As a result of their FH, the incidence of fatal or non-fatal myocardial infarction without treatment is about 50% by the age of 50 years in men and about 30% by the age of 60 years in women. (NHSEI). If the mother has FH, it is likely that the baby has been born with high cholesterol.

The current recommendation is that statins should be discontinued three months before attempting to conceive, during pregnancy and lactation (Shala-Haskaj 2020). During this time the cholesterol

level may become significantly raised even if the woman is eating a healthy, balanced diet and keeping active (NHS).

It is reported by Holmsen et al that the FELIC study showed that hypercholesterolaemia in a woman during pregnancy increased the risk of atherosclerosis in the child (Napoli 1999). They state that animal studies have shown that statin treatment in pregnant and lactating mice has a cardioprotective effect not only in the pregnant mouse but also in the offspring (Elahi 2008, 213)

Cholesterol levels are normally increased by 40% during pregnancy and lactation in healthy women (Lawrence 2016). The cholesterol in breast milk is synthesized in the mammary gland and its concentration in breast milk ranges from 27 mg/dL in colostrum to 16 mg/dL in mature breast milk (>30 days post-partum) (Lawrence 2016). Cholesterol is essential for brain tissue development, myelination of nerves and as a base for many enzymes. The role of higher cholesterol in colostrum is unclear (Lawrence 2016) The amount of cholesterol in breast milk that would hypothetically remain when a mother is taking a statin, would still be much higher than that provided by standard artificial formulas (Holmsen 2017). Standard artificial formula does not vary in composition at all and lacks cholesterol itself although it contains other lipids. Lawrence comments that interest in lipids in human milk has increased after reports of advance development at 12 months (Lucas 1992), 8-10 years (Lucas 1994) and even at 18 years (Horwood 1998). For evidence-based information on artificial formula milk please see: <https://www.firststepsnutrition.org/>

Use of statins during breastfeeding

Although there are no current studies on the use of statins in breastfeeding (Hale) in one study, breast milk from a woman with the familial hypercholesterolaemia gene contained three times as much cholesterol as that of healthy control women (Tsang 1978). Holmsen et al hypothesise that if cholesterol levels are normalised by statin use, it is reasonable to assume that the lipid content of breast milk will also decrease to more normal levels but will still exceed those of standard infant formula which contains no cholesterol.

Botha et al (Botha 2018) retrospectively reviewed 39 pregnancies from a cohort of 20 genotypically confirmed female patients with Homozygous Familial Hypercholesterolaemia (HoFH). Twenty-five pregnancies were exposed to lipid-lowering therapy, of which 18 were exposed to statin therapy, just prior to or during the pregnancy. The infants of HoFH patients are obligate Heterozygous familial hypercholesterolemia (HeFH) and likely already affected by atherosclerotic lesions at birth. Twelve of the 20 patients chose to breastfeed (21 infants). "Most patients breastfed for three to six months"; three patients opted to breastfeed for nine months. However, only 6 patients (11 infants) continued to breastfeed despite restarting statin therapy after delivery, while three patients (four infants) chose not to breastfeed while on statin therapy. Botha concluded that for many females with HoFH, despite the high cardiovascular risk, pregnancy is not uncommon and that lipid lowering therapy, particularly statin therapy during pregnancy, appears to be safe for both mother and foetus. Infants had no developmental or school learning problems. Botha et al recommended "patients should discontinue all lipid lowering therapy a month prior to planned conception and reinstate lipid lowering therapy, statin plus ezetimibe, during the second trimester. This limits the possible teratogenic effect of statins during the first trimester while providing the mother with optimal care during a time of increased cholesterol production. Women's perceptions and preferences regarding statin use in pregnancy should also be considered when giving advice in these situations. It is interesting that no recommendation is made on breastfeeding despite the health outcomes associated with infant feeding (UNICEF).

Pan (1988) studied 11 lactating women who were not breastfeeding but taking 20 mg of pravastatin orally twice daily for 2.5 days. Serum and milk samples were taken and analysed for pravastatin and its active metabolite after the fifth dose. Peak milk levels averaged 3.9 mcg/L for pravastatin and 2.1 mcg/L for its metabolite. He suggested that negligible levels were excreted into breast milk, but that benefits, and risks should be carefully considered. Using the peak levels above, a fully breastfed infant

would receive a maximum of 900 ng/kg daily with this dosage or about 0.13% of the maternal weight-adjusted dosage (LactMed).

Lwin (2018) studied a mother with a 13-month child still being breastfed who was commenced on Rosuvastatin 20mg at bedtime. The peak milk concentration was 58.6 mcg/L occurred 17 hours after the dose. The authors calculated a daily infant dosage of 4.63 mcg/kg, which corresponded to a weight-adjusted 1.5% of the maternal dose. Breastfeeding was discontinued so no analysis of the baby's plasma levels, or outcome were available.

Schutte (2018) published a case report of a woman with familial hypercholesterolemia who was started on rosuvastatin 40 mg daily on day 33 postpartum. Levels of the drug were measured up to day 80. All concentrations of the drug were in the range of 21 to 22 mcg/L. Despite the fact that samples were provided for 80 days there is no mention of the effect on the baby.

Pharmacokinetics of lipid lowering drugs

	Oral bioavailability %	Plasma protein binding %	Theoretical dose	Relative infant dose
Simvastatin	<5%	>95		
Atorvastatin	12 to 30	>98		
Pravastatin	17	50	0.001mg/Kg/d	0.15%
Rosuvastatin	20	88	0.003 mg/Kg/d	0.5 - 1 %
Ezetimibe	35 to 60	92-100		

The pharmacokinetic data (high percentage of protein binding, extensive first pass metabolism and low oral bioavailability) make it highly unlikely that significant quantities of statins will pass into breast milk (Elactancia). Data taken from Lennernäs 2003, NCBI StatPerls, Krishna 2009, Elactancia, Hale)

However, lack of passage into breastmilk has not been proven in research. Standard Infant formula milk contains no cholesterol but does contain other lipids.

Statins are grouped into low, medium, and high intensity according to the percentage reduction in low-density lipoprotein cholesterol (NICE CG 181 2016):

- a 20% to 30% reduction is low intensity e.g., pravastatin
- a 31% to 40% reduction is medium intensity
- a reduction of more than 40% is high intensity e.g., simvastatin 80mg, atorvastatin >20mg, rosuvastatin > 10mg.

MHRA (2014): there is an increased risk of myopathy associated with high dose (80 mg) simvastatin.

Adverse events from any drug on the breastfed baby should be reported to MHRA on a yellow card <https://yellowcard.mhra.gov.uk/>

New Drug Treatments

There are 2 new drugs used in specialist circumstances. However, there is no long-term data on the safety of using these new medications in breastfeeding mothers.

- Inclisiran (NICE TA 733)- Oral bioavailability 0-low, molecular weight 17,285 (EMA 2020)
- Evolocumab (NICE TA 394)- Oral bioavailability 0-low, molecular weight 144,000 (Hale, LactMed)

The two drugs discussed below, have large molecular weights and the amount in breast milk is expected to be exceptionally low. However, there is no long-term data on the safety of using these new medications in breastfeeding mothers.

Inclisiran for treating primary hypercholesterolaemia or mixed dyslipidaemia NICE [TA733]

Published: 06 October 2021.

There is no information available at all for the use of inclisiran whilst breastfeeding in humans. However, pharmacokinetics show: plasma concentrations reach undetectable levels within 48 hours of subcutaneous injection. Inclisiran is 87% protein bound with a half-life of approximately 9 hours. No accumulation occurs with multiple dosing (regimen of initial dose, followed by dose after 3 months, then every 6 months) ([emc 2021](#)). It is also a very large molecule (molecular weight (17,285Da) ([EMA2020](#)). Oral bioavailability is low.

Based on these pharmacokinetic factors, the passage of inclisiran into breastmilk would be expected to be low. Infants exposed to inclisiran via breastmilk should be monitored for any side effects especially poor feeding/ weight gain.

Evolocumab for treating primary hypercholesterolaemia and mixed dyslipidaemia NICE [TA394]

Published: 22 June 2016.

Evolocumab is a monoclonal antibody which are very large protein molecules, it has a molecular weight of 144,000Da (Hale) which suggests the amount transferred into breastmilk is likely to be very low. Oral bioavailability is low.

Based on these pharmacokinetic factors, the passage of Evolocumab into breastmilk would be expected to be low. Infants exposed to evolocumab via breastmilk should be monitored for diarrhoea and poor feeding/poor weight gain.

References

- Botha TC, Pilcher GJ, Wolmarans K, Blom DJ, Raal FJ. Statins and other lipid-lowering therapy and pregnancy outcomes in Homozygous Familial Hypercholesterolaemia: A retrospective review of 39 pregnancies. *Atherosclerosis*. 2018 Oct; 277:502-507. <https://pubmed.ncbi.nlm.nih.gov/30270091/>
- Drugs and Lactation Database (LactMed) <https://www.ncbi.nlm.nih.gov/books/NBK501922/>
- Elactancia database <https://www.e-lactancia.org/breastfeeding>
- Elahi 2008) MM, Cagampang FR, Anthony FW et al. Statin treatment in hypercholesterolemic pregnant mice reduces cardiovascular risk factors in their offspring. *Hypertension* 2008; 51: 939 – 44.
- Elahi MM, Cagampang FR, Ohri SK et al. Long-term statin administration to dams on high-fat diet protects not only them but also their offspring from cardiovascular risk. *Ann Nutr Metab* 2013; 62: 250 – 6).
- Hale TW *Medications and Mothers Milk* – online access HalesMeds.com
- HEARTUK What is Familial hypercholesterolaemia (FH) <https://www.heartuk.org.uk/cholesterol/what-is-fh>
- Holmsen ST, Bakkebø T, Seferowicz M, Retterstøl K. Statins and breastfeeding in familial hypercholesterolaemia <https://tidsskriftet.no/en/2017/05/commentary-and-debate/statins-and-breastfeeding-familial-hypercholesterolaemia>
- Horwood LJ, Fergusson DM. Breastfeeding and later cognitive and academic outcomes. *Pediatrics*. 1998 Jan;101(1): E9. doi: 10.1542/peds.101.1. e9. PMID: 9417173.
- Krishna R, Garg A, Jin B, et al. Assessment of a pharmacokinetic and pharmacodynamic interaction between simvastatin and anacetrapib, a potent cholesteryl ester transfer protein (CETP) inhibitor, in healthy subjects. *Br J Clin Pharmacol*. 2009;67(5):520-526. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2686068/> <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2686068/>
- LactMed Drugs and Lactation Database <https://www.ncbi.nlm.nih.gov/books/NBK501922/>

- L.L. Hui, Man Ki Kwok, E. Anthony S. Nelson et al (2019). Breastfeeding in Infancy and Lipid Profile in Adolescence. *Pediatrics*, Volume 143, Issue 5.
- Lawrence RA, Lawrence RM. Breastfeeding. A guide for the medical profession. Eighth Edition. Philadelphia: Elsevier; 2016
- Lennernäs H. Clinical pharmacokinetics of atorvastatin. *Clin Pharmacokinet*. 2003;42(13):1141-60.
- Lucas A, Morley R, Cole TJ, Gore SM. A randomised multicentre study of human milk versus formula and later development in preterm infants. *Arch Dis Child Fetal Neonatal Ed*. 1994;70(2): F141-F146. doi:10.1136/fn.70.2. f141
- Lucas A, Morley R, Cole TJ, Lister G, Leeson-Payne C. Breast milk and subsequent intelligence quotient in children born preterm. *Lancet*. 1992 Feb 1;339(8788):261-4.
- Lwin EMP, Leggett C, Ritchie U, et al. Transfer of rosuvastatin into breast milk: Liquid chromatography-mass spectrometry methodology and clinical recommendations. *Drug Des Devel Ther*. 2018; 12:3645–51.
- Napoli C, Glass CK, Witztum JL et al. Influence of maternal hypercholesterolaemia during pregnancy on progression of early atherosclerotic lesions in childhood: Fate of Early Lesions in Children (FELIC) study. *Lancet* 1999; 354: 1234 – 41
- NCBI StatPerls Pravastatin <https://www.ncbi.nlm.nih.gov/books/NBK551621/>
- Nguyen, B, Gale, J, Nassar, N, et al (2019). Breastfeeding and cardiovascular disease hospitalization and mortality in parous women: evidence from a large Australian cohort study. *Journal of the American Heart Association*, doi.org/10.1161/JAHA.118.011056
- NHS High Cholesterol <https://www.nhs.uk/conditions/high-cholesterol>
- NHS Lower your cholesterol <https://www.nhs.uk/live-well/healthy-body/lower-your-cholesterol/>
- NHS What is high cholesterol <https://www.nhs.uk/conditions/high-cholesterol>
- Owen CG et al (2002). Infant Feeding and Blood Cholesterol: A Study in Adolescents and a Systematic Review. *Pediatrics* 110: 597-608.
- Pan H, Fleiss P, Moore L, et al. Excretion of pravastatin, an HMG CoA reductase inhibitor, in breast milk of lactating women. *J Clin Pharmacol*. 1988; 28:942.
- Renfrew MJ, Pokhrel S, Quigley M, McCormick F, Fox-Rushby J, Dodds R, Duffy S, Trueman P, William A. Preventing disease and saving resources: the potential contribution of increasing breastfeeding rates in the UK 2012
- Schutte AE, Symington EA, du Preez JL. Rosuvastatin is transferred into human breast milk: A case report. *Am J Med*. 2013;126: e7–8
- Schwarz EB Ray RM, Steube AM et al. Duration of lactation and risk factors for maternal cardiovascular disease. *Obstet Gynecol* 2009 May; 113:974
- Shala-Haskaj P, Krähenmann F, Schmidt D. CME: Familiäre Hypercholesterinämie – Behandlung mit Statinen in der Schwangerschaft und Stillzeit [CME: Familial Hypercholesterolemia - Statin Treatment during Pregnancy and Breastfeeding]. *Praxis (Bern 1994)*. 2020 Apr;109(6):405-410. German
- Singhal A et al (2004). Breastmilk feeding and lipoprotein profile in adolescents born preterm: follow-up of a prospective randomised study. *Lancet* 363: 1571-78 <https://pubmed.ncbi.nlm.nih.gov/15145629/>
- Tsang RC, Glueck CJ, McLain C et al. Pregnancy, parturition, and lactation in Familial Homozygous Hypercholesterolemia. *Metabolism* 1978; 27: 823 – 9
- UNICEF Baby Friendly Initiative: the impact that breastfeeding can have on maternal cardiovascular health. <https://www.unicef.org.uk/babyfriendly/news-and-research/baby-friendly-research/maternal-health-research/maternal-health-research-heart-disease/>
- WHO Infant and young child feeding June 2021 <https://www.who.int/news-room/fact-sheets/detail/infant-and-young-child-feeding>

- [NHS England and NHS Improvement London \(NHSEI\) , Familial Hypercholesterolemia \(FH\)](https://www.england.nhs.uk/london/london-clinical-networks/our-networks/cardiac/familial-hypercholesterolaemia/)
- <https://www.england.nhs.uk/london/london-clinical-networks/our-networks/cardiac/familial-hypercholesterolaemia/>
- [NICE CG 181 \(2016\) Cardiovascular disease: risk assessment and reduction, including lipid modification](https://www.nice.org.uk/guidance/CG181)
- [MHRA Simvastatin: increased risk of myopathy at high dose \(80 mg\)](https://www.gov.uk/drug-safety-update/simvastatin-increased-risk-of-myopathy-at-high-dose-80-mg)
- <https://www.gov.uk/drug-safety-update/simvastatin-increased-risk-of-myopathy-at-high-dose-80-mg>
- NICE TAA 733 Inclisiran for treating primary hypercholesterolaemia or mixed dyslipidaemia 2021
- [Leqvio 284 mg solution for injection in prefilled syringe - Summary of Product Characteristics \(SmPC\) - \(emc\) \(medicines.org.uk\)](https://www.ema.europa.eu/en/documents/summary-product-characteristics/leqvio-284-mg-solution-for-injection-in-prefilled-syringe-summary-product-characteristics-sm-PC-epar-public_en.pdf)
- NICE TAa 394 Evolocumab for treating primary hypercholesterolaemia and mixed dyslipidaemia 2016
- [Repatha SureClick - Summary of Product Characteristics \(SmPC\) - \(emc\) \(medicines.org.uk\)](https://www.ema.europa.eu/en/documents/summary-product-characteristics/repatha-sureclick-summary-product-characteristics-sm-PC-epar-public_en.pdf)
- [Leqvio - 5333 - EPAR \(europa.eu\)](https://www.ema.europa.eu/en/medicines/human/EPAR/leqvio/leqvio-5333)
- [High cholesterol - Cholesterol levels - NHS \(www.nhs.uk\)](https://www.nhs.uk/conditions/high-cholesterol/)

The authors would like to express their gratitude to Dr James Akre and Prof Anders Hakansson for their support in the preparation of this information and to Kirstin Worsley, Supporter and Tutor The Breastfeeding Network as well as the other pharmacists of the Breastfeeding Network Drugs in Breastmilk Information Service team for their input.

Dr Wendy Jones MRPharmS MBE, Mrs S. Morris MRPharmS. Mrs A Da Costa MRPharmS